


A. **Spherical Coordinates.** There is another coordinate system that is excellent for objects that are symmetric about the origin, like spheres centered at the origin.

These are the spherical coordinates  $\rho$ ,  $\phi$ , and  $\theta$ . The spherical coordinate  $\theta$  is the same as the cylindrical coordinate. Let's explore in [Desmos](#).  $\rho$  is read "roe" and  $\phi$  is read "fee".




Every point can be described using coordinates in the range:

$$\rho \geq 0$$

$$0 \leq \theta \leq 2\pi$$


$$0 \leq \phi \leq \pi$$

For example we sketch the surface  $\mathcal{S}$  defined by  $\rho = 1, 0 \leq \phi \leq \frac{\pi}{2}, 0 \leq \theta \leq \frac{\pi}{2}$ .



On a sphere, the coordinates  $\phi$  and  $\theta$  are analogous to latitude and longitude.

B. **Spherical Conversion.** We develop algebraic relationships between spherical coordinates and the other coordinate systems.



**[Spherical ↔ Cylindrical] and [Spherical ↔ Standard]**

$r$  in terms of spherical coordinates:

$x$  in terms of spherical coordinates:

$y$  in terms of spherical coordinates:

$z$  in terms of spherical coordinates:


relation between  $\rho$  and standard/cylindrical coordinates:

relation between  $\phi$  and standard/cylindrical coordinates:


**Example 1.** Convert the cylindrical equation:

$$z = \sqrt{3}r$$

into a spherical equation, and sketch this surface assuming  $r \geq 0$ .



**C. Spherical Integration.** Now we will discuss integration in spherical coordinates. Imagine the infinitesimal bit of volume  $dV$  obtained from the point with spherical coordinates  $\rho, \phi, \theta$  by increasing  $\rho$  by  $d\rho$ ,  $\phi$  by  $d\phi$ , and  $\theta$  by  $d\theta$ .



The goal in this process is to try to understand how small changes  $d\rho$ ,  $d\phi$ , and  $d\theta$  are related to small changes  $dV$  in volume. This  $dV$  is exactly what appears in triple-integration.

For example, we know that small changes  $dr$ ,  $d\theta$ ,  $dz$  effect the change  $dV = r dr d\theta dz$  in volume.


In spherical coordinates:

$$dV =$$

**Example 2.** Use spherical coordinates to derive the volume of a radius  $R$  sphere:

$$x^2 + y^2 + z^2 \leq R^2$$

To describe a solid sphere of radius  $R$  centered at the origin, the section of the sphere in each vertical half-plane at angle  $\theta$  from  $0$  to  $2\pi$  is traced out by rotating the line segment from  $(0, 0, 0)$  to  $(0, 0, R)$  down from angle  $0$  to angle  $\pi$ .



**Example 3.** Let  $E$  be the region outside the cone  $z = \sqrt{x^2 + y^2}$  and inside the upper hemisphere of the sphere  $x^2 + y^2 + z^2 \leq 1$ . Find:

$$\iiint_E \frac{1}{\sqrt{x^2 + y^2 + z^2}} dV$$

The upper hemisphere of this sphere, which is centered at the origin, consists of the half of the sphere above the  $xy$ -plane.

